



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>A61K 31/00</b>	<b>A2</b>	(11) International Publication Number: <b>WO 00/64424</b> (43) International Publication Date: 2 November 2000 (02.11.00)
<p>(21) International Application Number: PCT/GB00/01564</p> <p>(22) International Filing Date: 20 April 2000 (20.04.00)</p> <p>(30) Priority Data: 9909135.7 22 April 1999 (22.04.99) GB</p> <p>(71) Applicant (for all designated States except US): UNIVERSITY OF WALES COLLEGE OF MEDICINE [GB/GB]; Research and Commercial Office, Heath Park, Cardiff CF4 4XN (GB).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): DORMER, Robert, Leslie [GB/GB]; University of Wales College of Medicine, Dept. of Medical Biochemistry, Heath Park, Cardiff CF4 4XN (GB). MCPHERSON, Margaret, Ann. [GB/GB]; University of Wales College of Medicine, Dept. of Medical Biochemistry, Heath Park, Cardiff CF4 4XN (GB).</p> <p>(74) Agent: BLAKEY, Alison, Jane; SmithKline Beecham, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i></p>
<p>(54) Title: CYSTIC FIBROSIS MEDICAMENTS</p> <p>(57) Abstract</p> <p>The use of a type V cyclic nucleotide phosphodiesterase inhibitor for the preparation of a medicament for the treatment of cystic fibrosis.</p>		

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CJ	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

### CYSTIC FIBROSIS MEDICAMENTS

The present invention relates to the use of a type V cyclic nucleotide phosphodiesterase inhibitor for the preparation of a medicament for the treatment of cystic fibrosis. In particular, it relates to the use of specific inhibitors for the preparation of a medicament for the treatment of cystic fibrosis.

Cystic fibrosis (CF) is an autosomal recessive disease characterised by disturbances in ion transport and viscous epithelial mucous secretions. The CF gene protein, CFTR acts as a Cl-channel and is also a key regulator of protein secretion.

CF is caused by mutations in the CF gene, which includes the cystic fibrosis transmembrane conductance regulator protein (CFTR).

WO-A-98/19679 discloses the use of 8-cyclopentyl theophylline (CPT) for the preparation of medicaments for the treatment of cystic fibrosis.

It has now been found that a type V cyclic nucleotide phosphodiesterase (PDE) inhibitor corrects the defective mucin secretory response to the  $\beta$ -agonist isoproterenol in CFTR antibody inhibited submandibular acini. Thus, a type V PDE inhibitor corrects defective CFTR function leading to rational drug treatment for cystic fibrosis.

A type V PDE inhibitor is as effective as IBMX or cpt-cyclic AMP. However, in contrast to IBMX and a selective type IV PDE inhibitor, the type V inhibitor has no effect on cyclic AMP levels and does not stimulate wild type mucin secretion. The data show that correction does not correlate with increase in cyclic nucleotide levels.


According to a first aspect of the present invention, there is provided the use of a type V cyclic nucleotide phosphodiesterase inhibitor in the preparation of a medicament for the treatment of cystic fibrosis.

Suitable type V cyclic nucleotide PDE inhibitors for use in the invention include those described in EP 347146, EP 351058, EP 352960 and in J. Med. Chem., 1993, 36(10), 1387-92.


The type V cyclic nucleotide PDE inhibitor is preferably a selective type V cyclic nucleotide PDE inhibitor.

Compounds can be identified as type V cyclic nucleotide PDE inhibitors using methods known to those skilled in the art, for example as described in EP 293063.

According to a second aspect of the present invention, there is provided the use of a compound of formula (I),




wherein A is a five- or six- membered ring containing one or more N atoms,  
X is a substituent of formula (II),




(II)

- wherein  $R_1$  represents  $C_{1-4}$  alkyl, and  
 X is positioned at either or both of the 2-position and/or the 8- position in (I), in the  
 5 preparation of a medicament for the treatment of cystic fibrosis.


Preferably, ring A in the compound of formula (I) is of the formula:



- wherein \* denotes the points of attachment of the ring, more preferably ring A is  
 10 of the formula:



Alternatively, ring A may be of the formula:




15

wherein  $R_2$  represents hydrogen or  $C_{1-4}$  alkyl and \* denotes the points of attachment of the ring.

Alternatively, ring A may be of the formula:

20




wherein \* denotes the points of attachment of the ring.


In particular, the substituent X is preferably only positioned at the 2-position, i.e. on the 4-pyrimidinone ring, in the compounds of formula (I).

25

According to a further aspect of the present invention, there is provided the use of a compound of formula (III):




wherein A is a five- or six- membered ring containing one or more N atoms,  
X is a substituent of formula (II):



wherein R<sub>1</sub> represents C<sub>1-4</sub> alkyl, and

X is positioned at either or both of the 2-position and/or the 8- position, in the preparation  
of a medicament for the treatment of cystic fibrosis.

Preferably, ring A of compound (III) has the following formula:



wherein \* denotes the points of attachment of the ring.

In particular, the substituent X is preferably only positioned at the 8-position, i.e.  
on ring A, in the compounds of formula (III).

The compounds of formulae (I) and (III) may be prepared as described in EP  
347146, EP 351058, EP 352960 and J. Med. Chem., 1993, 36(10), 1387-92.

In order to use type V cyclic nucleotide PDE inhibitors in therapy, they will  
normally be formulated into a pharmaceutical composition in accordance with standard  
pharmaceutical practice.

The type V cyclic nucleotide PDE inhibitors may be presented in a form suitable  
for oral administration in a pharmaceutical vehicle convenient for that administrative  
route. Thus, for example, the medicament may be presented as tablets, capsules,  
ingestible liquid or a powder preparation. Such formulations can include  
pharmaceutically acceptable carriers known to those skilled in the art. Formulations  
suitable for oral administration further include lozenges, pastilles, aerosols and  
mouthwashes. The type V cyclic nucleotide PDE inhibitors may also be administered by  
inhalation, that is by intranasal and oral inhalation administration. Formulations suitable  
for inhalation include powders where the carrier is a solid, and where the carrier is a  
liquid, the formulation can be administered as a nasal spray or aerosol, or as drops for  
example.

In yet another alternative, the type V cyclic nucleotide PDE inhibitors may be  
presented in a formulation suitable for parenteral intravenous administration, such as

aqueous or non-aqueous sterile injectable solutions. Such solutions may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient. Alternatively, the formulation for parenteral administration may be presented as an aqueous or non-aqueous sterile suspension which may include suspending agents and thickening agents.

It will also be recognized by one of skill in the art that the optimal quantity and spacing of individual dosages of the inhibitors will be determined by the nature and extent of the condition being treated, the form, route and site of administration, and the particular patient being treated, and that such optimums can be determined by conventional techniques. It will also be appreciated by one of skill in the art that the optimal course of treatment, i.e., the number of doses of the inhibitors given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

According to a further aspect of the present invention, there is provided a method of treating cystic fibrosis comprising administering a type V cyclic nucleotide phosphodiesterase inhibitor to a patient in need thereof. The type V cyclic nucleotide PDE inhibitor may be a compound of formula (I) or formula (III) as mentioned above.

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

Whilst the invention has been described above, it extends to any inventive combination of the features set out above or in the following description. The invention will now be described, by way of example, with reference to the following drawing and examples.

Figure 1 shows the correction of CFTR antibody-inhibited  $\beta$ -adrenergic stimulation of mucin secretion by PDE inhibitors.

## Examples

### Methods

#### Production of anti-peptide CFTR antibodies

A peptide consisting of 14 amino acids (524-537), residing in the first nucleotide binding domain (NBD) region of CFTR, was synthesized and coupled to keyhole limpet haemocyanin (KLH) (10mg peptide/8mg KLH) by Cambridge Research Biochemicals Ltd, Northwich, Cheshire. The peptide sequence was searched for in the Swissprot database (release of 21/3/92, Daresbury Lab., Seqnet facility) and only CFTR was found to contain a perfect match. Antisera were prepared as described (Lloyd Mills et al, (1992) Biochem. Biophys. Res. Commun., 188, 1146-1152 and Pereira et al, (1998) Brit. J. Pharmacol., 125, 697-704) and affinity-purified using KLH or peptide coupled to CH-Sepharose 4B (Pharmacia). The IgG content of antisera was estimated either following precipitation with 24%  $\text{Na}_2\text{SO}_4$  for 2h at room temp. (KLH-Sepharose purified) or directly (peptide-Sepharose purified), using the BioRad protein assay kit.

### Isolation of rat submandibular acini, incorporation of antibodies into intact acini and measurement of mucin secretion


Procedures were carried out as previously described (Lloyd Mills et al, Biochem. Biophys. Res. Commun., 1992, 188, 1146-1152, and Pereira et al, Br. J. Pharmacol., 1998, 125, 697-681). Briefly, acini were pulse-chase labelled with [<sup>3</sup>H]-glucosamine and suspended in TES-buffered saline (10mM TES (N-Tris-(hydroxymethyl)-methyl-2-aminoethanesulphonic acid), pH 7.4 containing 143mM NaCl, 4.7mM KCl, 1.1mM MgCl<sub>2</sub>, 1mg/ml bovine serum albumin BSA). An equal volume of either 10mM TES, pH 7.4 (swollen) or TES-buffered saline (unswollen), each containing 5mM ATP and antibody (1mg/ml) was added for 1.5 min at room temperature, followed by washing and resuspension in Krebs-Henseleit bicarbonate (KHB) buffer containing 20mg/ml BSA. Following a 15 min recovery incubation at 37°C in KHB buffer, acini were washed and incubated under experimental conditions at 37°C. Isoproterenol and 3-isobutyl-1-methylxanthine (IBMX) were dissolved directly into KHB medium at the concentrations used; the selective type III-V PDE inhibitors were dissolved initially in DMSO and diluted to give a final concentration of <1% DMSO in the incubations. An equivalent amount of DMSO was added to control incubations. Type V PDE inhibitor was the least soluble; hence a concentration of 0.2mM, rather than 1mM was used. [<sup>3</sup>H]-labelled mucins, released into the medium at zero time and after 30 min, were acid-precipitated and their radioactivity measured as described by Lloyd Mills et al, Biochem. Biophys. Res. Commun., 1992, 188, 1146-1152, and Pereira et al, Br. J. Pharmacol., 1998, 125, 697-681. Protein content of cell pellets was determined using the BioRad protein assay kit and mucin release over 30 min expressed as dpm/mg protein or as % basal secretion to take account of variation in unstimulated mucin release between experiments.

### Measurement of cyclic AMP and cyclic GMP content


Aliquots of acini suspensions (0.5ml) were added to an equal volume of ice cold trichloroacetic acid (20%), extracted and assayed using specific radioimmunoassay kits for cyclic AMP and cyclic GMP (Amersham), as described by Lloyd Mills et al, Biochem. Biophys. Res. Commun., 1992, 188, 1146-1152.

### Results


The mechanism of action of IBMX in correcting defective CFTR function was investigated using (R,S)-4,5-dihydro-6-[4-(1,4-dihydro-4-oxopyridin-1-yl)phenyl]-5-methyl-3(2H)-pyridazinone a selective type III (cyclic AMP activated, cyclic GMP inhibited) PDE inhibitor, 4-(3-cyclopentyloxy-4-methoxyphenyl)-2-pyrrolidinone a type IV (cyclic AMP activated) PDE inhibitor, and 1,7-dihydro-2-(2-propoxyphenyl)-6H-purin-6-one a type V (cyclic GMP activated) PDE inhibitors. The structure of these compounds is shown below:




IBMX



Type III



Type IV



Type V

Table 1 shows the actions of maximally effective concentrations of isoproterenol (IP), IBMX and the selective cyclic nucleotide PDE inhibitors on mucin secretion and cyclic AMP rise from rat submandibular acini. Mucin secretion was measured at 30 min; cyclic AMP at 5 min, when maximum stimulation is seen. Isoproterenol and IBMX were the most potent stimulators of mucin secretion, giving a similar maximum stimulation; however isoproterenol increased cyclic AMP to a much greater extent than IBMX. The type IV PDE inhibitor gave the same cyclic AMP rise as isoproterenol but was much less effective in stimulating mucin secretion. The type III and type V PDE inhibitors did not significantly increase cyclic AMP or mucin secretion. The data indicate a lack of correlation between magnitude of increase in cyclic AMP and stimulation of mucin secretion and are in accord with previous results (Bradbury et al, Biochem. Biophys. Res. Commun., 1998, 161, 661-671 and Lloyd Mills et al, FEBS Lett., 1991, 298, 141-144) showing dissociation of cyclic AMP rise and mucin secretion which suggest that cyclic AMP is not the sole intracellular messenger involved.

Figure 1 shows the actions of the selective cyclic nucleotide PDE inhibitors on the defective isoproterenol-stimulated mucin secretion in rat submandibular cells containing CFTR antibody (approx. 1mg IgG/ml) which had been introduced as described in the



Methods by hypotonic swelling. The actions of the PDE inhibitors in increasing the CFTR antibody-inhibited response was compared to that of cpt-cyclic AMP, which has been shown to restore secretory responsiveness to the same extent as IBMX (1mM), giving approx. 75% of the secretion response seen in cells containing an equivalent amount of non-immune IgG. The data show that type IV and type V PDE inhibitors are as effective as cpt-cAMP in correcting the defective B-adrenergic secretory response, whereas the type III PDE inhibitor was ineffective. Referring to Figure 1, rat submandibular acini were pulse-chase labelled with  $^3\text{H}$ -glucosamine and swollen in the presence of CFTR antibody as described in the Methods. Mucin secretion in the presence of isoproterenol, (Iso-10 $\mu\text{M}$ ) and PDE inhibitors (PDE I Types III and IV at 1mM; Type V at 0.2mM) or cpt-cyclic AMP (1mM) as shown, was measured after 30 min. Results are mean  $\pm$  SEM for at least 4 experiments and are expressed as % basal. \*,  $p < 0.05$ , \*\*,  $p < 0.01$  for difference from secretion in the presence of isoproterenol alone as assessed by Student's t-test.

IBMX potentiates the cyclic AMP rise in the presence of isoproterenol, giving excessive increase in cyclic AMP above the levels induced physiologically by B-adrenergic stimulation, although it does not increase mucin secretion above the maximum induced by isoproterenol. The actions of the selective cyclic nucleotide PDE inhibitors on isoproterenol stimulated mucin secretion and cyclic AMP rise have been compared to that of IBMX as shown in Table 2. Previous results have shown that the cyclic AMP response of non-swollen cells or cells swollen in the presence of either non-immune IgG or CFTR antibody are not different. It is noteworthy that like IBMX, none of the selective PDE inhibitors significantly increased mucin secretion above the maximum response induced by isoproterenol alone, even through IBMX, type III and type IV inhibitors massively potentiated the isoproterenol induced cyclic AMP rise. Table 2 also shows that the type III PDE inhibitor was as effective as IBMX in potentiating the cyclic AMP rise induced by isoproterenol alone and the type IV inhibitor was even more effective. However, the Type V PDE inhibitor did not potentiate the isoproterenol induced cyclic AMP response. This raised the question as to the role of cyclic GMP in the correction of CFTR function by the Type V PDE inhibitor. Table 3 shows that isoproterenol alone caused a very small increase in cyclic GMP, which was further increased in the presence of IBMX or the selective Type V PDE inhibitor. Type III and type IV PDE inhibitors did not change cyclic GMP in the presence or absence of isoproterenol (Table 3).

Further selective type V PDE inhibitors as shown in Table 4 were also tested for their actions in correcting CFTR function. Table 5 shows their potency of correction compared to 1mM cpt-cyclic AMP.

The mechanism of action of the selective PDE inhibitors in correcting defective CFTR function did not correlate with their ability to stimulate mucin secretion nor to cause excessive increase in cyclic AMP. Thus, the type IV PDE inhibitor stimulated mucin secretion, gave excessive increase in cyclic AMP and corrected CFTR activity whereas the type III PDE inhibitor did not stimulate mucin secretion, gave excessive

increase in cyclic AMP but did not correct CFTR activity. The type V PDE inhibitor corrected CFTR activity without stimulating mucin secretion or cyclic AMP.

- 5 Table 1. Actions of cyclic nucleotide PDE inhibitors on mucin secretion and cyclic AMP levels in rat submandibular acini.

Additions	Mucin Secretion (% basal)	Cyclic AMP (pmol/mg protein)
No addition	100±9	9.4±2.7
Isoproterenol (10µM)	**619±101	"160.3±23.1
IBMX (1mM)	**578±84	'27.9±4.9
Type III (1mM)	139±42	11.8±1.7
Type IV (1mM)	*191±20	'168±36.6
Type V (0.2mM)	122±35	4.9±0.6

- 10 Mucin release was measured at 30 min and cyclic AMP at 5 min as described in the Methods, in the presence or absence of agonists. Mucin release is expressed as % basal, due to the variation in basal between experiments. Results are means ± SEM for at least three experiments. Significance of differences was assessed by Students t-test: \*, p<0.002; \*\*, p<0.001 for difference from basal mucin secretion; ', p<0.05; ", p<0.01 for difference from basal cyclic AMP.

- 15 Table 2. Actions of cyclic nucleotide PDE inhibitors on isoproterenol-stimulated mucin secretion and cyclic AMP levels in rat submandibular acini.

	Mucin Secretion (% basal)	Cyclic AMP (pmol/mg protein)
No addition	100± 7 (6)	7.9±1.9 (5)
Isoproterenol (10µM)	785±212 (6)	373.0±84.0 (6)
Isoproterenol (10µM)	924±290 (6)	*1252.9±129.6 (6)
+ IBMX (1mM)		
Isoproterenol (10µM)	692±238 (6)	*1009.6±91.5 (6)
+ Type III (1mM)		
Isoproterenol (10µM)	708±220 (6)	**1908.7±150.8 (6)
+ Type IV (1mM)		
Isoproterenol (10µM)	755±243 (6)	375.5±95.8 (6)
+ Type V (0.2mM)		

- 20 Mucin secretion was measured at 30 min and cyclic AMP at 5min as described in the Methods, under the conditions shown. Mucin secretion is expressed as % basal, due to the variation between experiments. Numbers in parentheses represent the number of observations. Significance of difference was assessed by Students unpaired t-test. \*, p<0.001; \*\*, p<0.0001 for difference from stimulation by isoproterenol alone.

Table 3. Actions of cyclic nucleotide PDE inhibitors on cyclic GMP levels in the presence or absence of isoproterenol.

	Cyclic GMP (fmol/mg protein)	
	No Isoproterenol	+ Isoproterenol (10 $\mu$ M)
No addition	39.2 $\pm$ 1.0 (6)	45.2 $\pm$ 1.6 (4)
IBMX (1mM)	48.1 $\pm$ 5.2 (8)	**55.6 $\pm$ 1.6 (8)
Type III (1mM)	39.3 $\pm$ 2.4 (3)	47.0 $\pm$ 1.8 (4)
Type IV (1mM)	40.1 $\pm$ 1.5 (3)	45.2 $\pm$ 2.0 (4)
Type V (0.2mM)	*47.3 $\pm$ 2.5 (5)	**64.3 $\pm$ 3.6 (5)

5





Cyclic GMP levels were measured as described in the Methods following 5 min incubation under the conditions shown. PDE inhibitor concentrations used were 1mM for IBMX, Types III and IV and 0.2mM for Type V. Results are mean  $\pm$  SEM for the number of experiments shown in parentheses. Significance of difference was assessed by

10

Students t-test. \*,  $p < 0.02$  for difference from basal (no isoproterenol); all conditions in the presence of isoproterenol were significantly different from basal ( $p < 0.05$ ); \*\*,  $p < 0.005$  for difference from stimulation by isoproterenol alone.

Table 4.

15

COMPOUND	STRUCTURE	PDE V IC <sub>50</sub> ( $\mu$ m)
a)		0.96
b)		0.55
c)		22
d)		34







e)		0.65
f)		4.5
g)		0.75
h)		48
i)		0.2
j)		1.3

Table 5. Actions of compounds on correction of CFTR function, wild type mucin secretion and cyclic AMP and cyclic GMP levels.

5

Compound	Degree of Correction compared to 1mM cAMP % cAMP response	Mucin secretion % basal	cyclic AMP % basal	cyclic GMP % basal
a)	100*	120	100	140
b)	100*	180*	600*	130
c)	100*	160*	250*	90
d)	100*	200*	600*	90
e)	50	170*	350*	130
f)	33	160*	100	110
g)	20	200*	160*	110

		180*	100	110
h)	0	170*	100	90
i)	0	160*	100	140*
j)	0			

\* indicates significant difference from basal.

5        The results show that a selective Type V cyclic nucleotide PDE inhibitor corrected defective  $\beta$ -adrenergic stimulation of mucin secretion induced by CFTR antibody. The Type V PDE inhibitor did not increase cyclic AMP levels, nor did it potentiate isoproterenol induced cyclic AMP rise. The data show that the CFTR mucin secretion defect can be corrected without increase in cyclic AMP and imply specificity in

10      correction of the CFTR defect.

A lack of correlation between magnitude of increase in cyclic AMP and stimulation of mucin secretion in submandibular cells was observed in that IP and IBMX maximally stimulated mucin secretion, but IBMX gave a much lower cAMP response and the type IV PDE inhibitor increased cyclic AMP to the same extent as IP, but gave

15      much less stimulated mucin secretion. The data are in accord with previous results showing dissociation between cAMP rise and mucin secretion, indicating that this is not the sole intracellular messenger involved. Stimulation of wild type mucin secretion was not a prerequisite for correction of CFTR activity, as shown by the lack of stimulation of mucin secretion by the active type V PDE inhibitor.

20      It is not clear whether the non selective PDE inhibitor IBMX and the type IV PDE inhibitor, which increase cyclic AMP and potentiate IP-induced cyclic AMP rise, correct defective CFTR function by an excessive increase in cyclic AMP. Results using the type III PDE inhibitor, showing the same isoproterenol induced cyclic AMP rise as IBMX but no restoration of CFTR activity, suggest that excessive increase in cyclic AMP alone is

25      not sufficient to correct defective CFTR function.

The finding that a Type V PDE inhibitor corrected CFTR function, raised the question as to whether cyclic GMP was involved. The predominant PDE in submandibular tissues is the type IV enzyme. In submandibular acini, isoproterenol increases cyclic GMP to a small extent, which is further increased in the presence of


30      IBMX or the type V PDE inhibitor. Thus the mechanism of correction by the type V PDE inhibitor may be mediated by cyclic GMP, but other actions such as direct interaction with CFTR are also possible, and the applicant is not to be bound thereby.

In accord with these findings, other structurally related analogues of the selective type V PDE inhibitor are active in correcting CFTR function (Table 5). Whilst the


35      applicant is not to be bound hereby, these compounds could be directly interacting with CFTR.

## CLAIMS

1. The use of a type V cyclic nucleotide phosphodiesterase inhibitor in the preparation of a medicament for the treatment of cystic fibrosis.
- 5 2. The use according to claim 1 wherein the type V cyclic nucleotide phosphodiesterase inhibitor is a selective type V cyclic nucleotide phosphodiesterase inhibitor.
- 10 3. The use of a compound of formula (I):




- wherein A is a five- or six- membered ring containing one or more N atoms,
- 15 X is a substituent of formula (II):




- wherein R<sub>1</sub> represents C<sub>1-4</sub> alkyl, and
- 20 X is positioned at either or both of the 2-position and/or the 8- position in (I), in the preparation of a medicament for the treatment of cystic fibrosis.

4. The use according to claim 3 wherein ring A is of the formula:



- 25 wherein \* denotes the points of attachment of the ring.


5. The use according to claim 3 wherein ring A is of the formula:



wherein  $R_2$  represents hydrogen or  $C_{1-4}$  alkyl and \* denotes the points of attachment of the ring.

5

6. The use according to claim 3 wherein ring A is of the formula:




wherein \* denotes the points of attachment of the ring.

10

7. The use according to claim 3 wherein the compound of formula (I) is 1,7-dihydro-2-(2-propoxyphenyl)-6H-purin-6-one.


8. The use of a compound of formula (III):

15



wherein A is a five- or six- membered ring containing one or more N atoms,  
X is a substituent of formula (II):


20



wherein  $R_1$  represents  $C_{1-4}$  alkyl, and  
X is positioned at either or both of the 2-position and/or the 8- position in (III),  
in the preparation of a medicament for the treatment of cystic fibrosis.

25

9. The use according to claim 8 wherein ring A is of the formula:



wherein \* denotes the points of attachment of the ring.

10. The use according to claim 8 or 9 wherein the substituent X is only positioned at  
5 the 2-position of ring A in a compound of formula (I).



FIGURE 1

